arXiv:0801.0470v1 [cond-mat.mes-hall] 3 Jan 2008

Spin-orbit interaction effect on transport of Dirac fermions in graphene

Kai-He Ding[1],* Guanghui Zhou[2],† and Zhen-Gang Zhu[3]
[1]*Department of Physics and Electronic Science, Changsha University of Science and Technology, Changsha, 410076, China*
[2]*Department of Physics, Hunan Normal University, 410081, China and*
International Center for Materials Physics, Chinese Academy of Sciences, Shenyang 110015, China and
[3]*Institut für Physik Martin-Luther-Universität Halle-Wittenberg Nanotechnikum-Weinberg,*
Heinrich-Damerow-Strasse 4 D - 06120 Halle (Saale), Germany

We study theoretically the quantum transport properties of the Dirac fermions with spin-orbit interactions (SOIs) in graphene by using the method of Schwinger proper time together with decomposition over Landau level poles and Kubo formula. The analytical expressions for both longitudinal and Hall conductivities are derived explicitly. It is found that, from some numerical examples, when the Rashba SOI is taken into account the Shubnikov-de Haas (SdH) oscillation peaks of the longitudinal conductivity versus the chemical potential are split, while the SdH oscillation of the longitudinal conductivity versus a external magnetic field exhibits a beating pattern. Furthermore, the Rashba SOI tends to suppress the quantum Hall effect in graphene.

PACS numbers: 73.50.-h,71.70.Ej,81.05.Uw

I. INTRODUCTION

Graphene has attracted a lot of attention because of its appealing properties[123456]. At low energy, owing to the specific band structure with the unique valley and neutrality separating the hole states from the electron states[7], graphene has led to the emergence of a paradigm of the relativistic condensed matter physics[89], where the relativistic quantum phenomena, some of which are unobservable in high energy physics, can be tested in the table-top experiment. The recent advances in fabrication technique have made it possible to produce graphite systems with a few layers or even a single monolayer of graphene[10111213], which propel graphene study into thriving high tide.

The relativistic feature of the graphene represents the anomalous physical phenomena, such as anomalous quantized Hall effect, absence of the weak localization and existence of the minimal conductivity[3]. In addition to dissipative transport also supercurrent transport has already been observed[4]. It is suggested that the graphene is a promising candidate for the spintronics and related applications due to the SOIs[1415]. In a recent paper, Kane and Mele have studied the effect of the SOIs in graphene[1617], and find that the spin Hall conductivity is quantized in the absence of a magnetic field because of a gap produced by the SOIs. Depending on the relative strength of intrinsic and Rashba SOIs, it is further shown that the spin Hall conductivity can be zero or nonzero. Kane and Mele have roughly estimated the SOI scale. Following this routine, Yao *et al.*[18] and Min *et al.*[19] have found that this estimation is too big, and they have given some explicit expressions of the SOIs in graphene.

In this work, we investigate the transport of Dirac fermions in graphene. The main purpose is focused on the effect of the SOIs on the transport on the basis of estimation of Yao *et al.* and Min *et al.* for the SOIs in graphene. Using the Schwinger proper-time method[20], decomposition over Landau level poles[2122] and Kubo formula[23], we obtain some analytical expressions for both longitudinal and Hall conductivities. It is found that when the Rashba SOI is considered, the longitudinal conductivity as a function of the chemical potential deviates the linear relation at zero magnetic field. For nonzero magnetic field, the SdH oscillations are observed and the oscillation peaks in the longitudinal conductivity versus the chemical potential are split when the Rashba SOI is applied, while the oscillation in the longitudinal conductivity versus the magnetic field exhibits a beating pattern. It is also shown that the Rashba SOI tends to suppress the quantum Hall effect in graphene.

The rest of this paper is outlined as follows: In Sec. II, The model of a single layer of graphite (graphene) with the SOIs is established. In Sec. III, we derive the analytical expressions for both the longitudinal conductivity and the Hall conductivity including the limits of these expressions at zero field. In Sec. IV, the corresponding numerical results and discussions are given. In Sec. V, a summery is presented. Finally, Some tedious algebra is included in the Appendix.

II. MODEL FORMALISM

The graphene is a flat monolayer of carbon atoms tightly packed into a honeymoon lattice. At low energy, it can be described by 2+1 dimensional relativistic field theory model. When the SOIs are included, the Lagrangian density of the system is given by

$$\mathcal{L} = \hbar v_F \overline{\Psi}(i\hat{D} + H_s)\Psi, \tag{1}$$

where $\Psi = (\Psi_K, \Psi_{k'})$ is the eight-component Dirac spinors with $\Psi_{K(K')} = (\Psi_{A\uparrow}, \Psi_{A\downarrow}, \Psi_{B\uparrow}, \Psi_{B\downarrow})$ which describes the spin-related Bloch states residing on the atoms of the A, B sublattice at momentum $K(K')$, $\hat{D} = \gamma^\mu(\partial_\mu - ieA_\mu)$ with γ^μ ($\mu = 0, 1, 2$) being 4×4 γ matrices[26], e is the electron charge, v_F is the Fermi velocity, the external magnetic field $\mathbf{B} = \nabla\times\mathbf{A}$ is applied perpendicular to the $x-y$ plane and the corresponding vector potential is taken in the symmetric gauge $\mathbf{A} = (-By/2, Bx/2)$. In Eq. (1), H_s describes the SOIs that read[1819]

$$H_s = \lambda_{SO}(1 - \gamma^0 s_z) + \lambda_R(i\gamma^1 s_y - \gamma^0\gamma^1 s_x), \tag{2}$$

where λ_{SO} is the intrinsic SOI parameter, λ_R is the Rashba SOI parameter, and $\mathbf{s}$ is the spin variable. For $B = 0$, the corresponding energy spectrum are given by

$$\varepsilon_1 = \pm\sqrt{\mathbf{k}^2 + (\lambda_R - \lambda_{SO})^2} + \lambda_R - \lambda_{SO},$$
$$\varepsilon_2 = \pm\sqrt{\mathbf{k}^2 + (\lambda_R + \lambda_{SO})^2} - \lambda_R - \lambda_{SO}. \quad (3)$$

For $\lambda_{SO} > \lambda_R > 0$, the system includes an energy gap of $2(\lambda_{SO} - \lambda_R)$. For $0 < \lambda_{SO} < \lambda_R$, the energy gap closes.

The Green's function of Dirac fermions described by the Lagrangian (1) in an external magnetic field can be expressed as[20,21,22]

$$G(x,y) = (i\hat{D} - H_s)_x\langle x|\frac{-i}{H_s^2 + \hat{D}^2 - i[\hat{D}, H_s]}|y\rangle. \quad (4)$$

Using the Schwinger proper time approach[20], we obtain

$$G(x,y) = \exp(ie\int_y^x A_\lambda dz^\lambda)\widetilde{G}(x-y), \quad (5)$$

$$\widetilde{G}(x) = \int_0^\infty ds\frac{e^{-\frac{i\pi}{4}}}{8(\pi s)^{3/2}}e^{-\frac{i}{4s}x_\nu C^{\gamma\mu}x_\mu}[\frac{1}{2s}\gamma^\mu C_{\mu\nu}x^\nu - \frac{1}{2}(e\gamma^1 Bx_2 - e\gamma^2 Bx_1) + \lambda_{SO}(1-\gamma^0 s_z)]\frac{eBs}{\sin(eBs)}e^{i(\frac{1}{2}e\sigma F - \Delta - B_\mu^2)s}, \quad (6)$$

where $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$, $C^{\mu\nu} = g^{\mu\nu} + \frac{(F^2)^{\mu\nu}}{B^2}[1 - eBs\cot(eBs)]$ with $g^{\mu\nu} = \mathrm{diag}(1,-1,-1)$, and

$$B_\mu = i2\lambda_{SO}\sigma^{\mu 0}s_z + 2\lambda_R(\sigma^{\mu 1}s_y - \delta_{\mu 0}\gamma^1 s_x - \delta_{\mu 1}\gamma^0 s_x), \quad (7)$$

$$\Delta = 2(\lambda_{SO}^2 + \lambda_R^2)(1 - \gamma^0 s_z) + 4\lambda_{SO}\lambda_R(i\gamma^1 s_y - \gamma^0\gamma^1 s_x) \quad (8)$$

with $\sigma^{\mu\nu} = \frac{i}{2}(\gamma^\mu\gamma^\nu - \gamma^\nu\gamma^\mu)$. It is clear that the symmetric gauge sets the factor $\exp(ie\int_y^x A_\lambda dz^\lambda)$=1 in Eq. (5). Therefore, Eq. (5) becomes the functions of the difference $x - y$ only. Using the expansions of the exponential function e^x, one can show that

$$\begin{aligned} e^{i(\frac{1}{2}e\sigma F - \Delta - B_\mu^2)s} &= e^{-i(10\lambda_{SO}^2 + 2\lambda_R^2)s}[\cos(\zeta s) + i\gamma^0 s_z\sin(\zeta s)]\{\cos(eBs) + \gamma^1\gamma^2\sin(eBs) + \frac{1}{2}(1-\gamma^0 s_z)[\cos(\xi s) - \cos(eBs)] \\ &- i\frac{12\lambda_{SO}\lambda_R\sin(\xi s)}{\xi}(i\gamma^1 s_y - \gamma^0\gamma^1 s_x) + \frac{1}{2}(1-\gamma^0 s_z)\gamma^1\gamma^2[\frac{eB}{\xi}\sin(\xi s) - \sin(eBs)]\}, \end{aligned} \quad (9)$$

where $\zeta = 2\lambda_{SO}^2 - 6\lambda_R^2$, $\xi = \sqrt{(24\lambda_{SO}\lambda_R)^2 + (eB)^2}$. Substituting Eq. (**??**) into Eq. (5), applying Fourier transform in the Matsubara representation and using the decomposition method over Landau level poles[21,22], we can derive the expression

$$G(i\omega_m, \mathbf{k}) = G_1(i\omega_m, \mathbf{k}) + G_2(i\omega_m, \mathbf{k}) + G_3(i\omega_m, \mathbf{k}) \quad (10)$$

after straightforward but somewhat complicated calculations. The expressions of $G_i(i\omega_m, \mathbf{k})$ (i=1,2,3) are very complicated and will be given in the appendix A. Whence, we can further obtain the retarded and advanced Green's functions by the analytic continuation $G^{(R)}(\omega + i0, \mathbf{k}) = G(i\omega_m \to \omega + i0, \mathbf{k})$ and $G^{(A)}(\omega - i0, \mathbf{k}) = G(i\omega_m \to \omega - i0, \mathbf{k})$. When considering the influence of impurities, it is assumed that the scattering rate Γ on impurity is described phenomenologically by a constant, and then the Green's functions acquire the form

$$G^{(R,A)}(\omega, \mathbf{k}) = G_1^{(R,A)}(\omega \pm i\Gamma, \mathbf{k}) + G_2^{(R,A)}(\omega \pm i\Gamma, \mathbf{k}) + G_3^{(R,A)}(\omega \pm i\Gamma, \mathbf{k}). \quad (11)$$

In general, the scattering rate Γ, which is defined by $\Gamma(\omega) = -Im\Sigma^R(\omega)$, is a frequency-dependent quantity. It needs to be determined self-consistently from the Schwinger-Dyson equations. The exact form of this equation actually depends on the impurity scattering fashion, such as short- or long-range scatterers. This kind of consideration have been made for graphene in Ref.[24]. But in this paper, we mainly focus on the SOI effect on transport, and neglect the exact form of in-

teractions between impurities and electrons.

III. ELECTRONIC CONDUCTIVITY

The Kubo formula concerning the frequency-dependent electrical conductivity as a linear response function to an external field can be written as[23]

$$\sigma_{ij}(\Omega) = \frac{\mathrm{Im}\Pi^R_{ij}(\Omega + i0)}{\Omega}, \tag{12}$$

where i, j are the component indexes of coordinates, and $\Pi^R_{ij}(\omega)$ is the retarded current-current correlation function obtained by analytical continuation of the Matsubara function

$$\Pi_{ij}(i\omega_n) = \frac{1}{V}\int_0^\beta d\tau e^{i\omega_n\tau}\langle T_\tau J_i(\tau)J_j(0)\rangle,\ \omega_n = 2\pi nT, \tag{13}$$

where V is the volume of the system, β is the inverse temperature, and $J_i(\tau) = \int d^2r\, j_i(\tau, \mathbf{r})$ with $j_i = -ev_F\overline{\Psi}(\tau, \mathbf{r})\gamma^i\Psi(\tau, \mathbf{r})$. Neglecting the impurity vertex corrections, the calculation of the conductivity reduces to evaluation of the bubble diagram. Then Eq. (12) can be rewritten as

$$\begin{aligned}\sigma_{ij}(\Omega) &= \frac{e^2v_F^2}{2\pi\Omega}\mathrm{Re}\int_{-\infty}^{\infty}d\omega\int\frac{d^2k}{(2\pi)^2}tr\{[n_F(\omega) - n_F(\omega+\Omega)][\gamma^iG^R(\omega+\Omega,\mathbf{k})\gamma^jG^A(\omega,\mathbf{k})]\\ &+ n_F(\omega+\Omega)\gamma^iG^A(\omega+\Omega,\mathbf{k})\gamma^jG^A(\omega,\mathbf{k}) - n_F(\omega)\gamma^iG^R(\omega+\Omega,\mathbf{k})\gamma^jG^R(\omega,\mathbf{k})]\},\end{aligned} \tag{14}$$

where $n_F(\omega)$ is the Fermi distribution function. Substituting Eq. (11) into Eq. (14), we can obtain the longitudinal conductivity

$$\sigma_{xx} = \frac{2e^2v_F^2|eB|}{\pi^2}\mathrm{Re}\int_{-\infty}^{\infty}d\omega\frac{1}{4T\cosh^2\frac{\beta(\omega-\mu)}{2}}A_L(\omega) \tag{15}$$

and the Hall conductivity

$$\sigma_{xy} = -\frac{2e^2v_F^2|eB|sgn(eB)}{\pi^2}\mathrm{Im}\int_{-\infty}^{\infty}d\omega\frac{1}{4T\cosh^2\frac{\beta(\omega-\mu)}{2}}A_H(\omega), \tag{16}$$

where all the quantities on the right-hand side are calculated in the Appendix B. Eqs. (15) and (16) establish the fundamental basis for investigating the SOI effect on the quantum transport properties of the Dirac fermions in graphene.

In the limit of zero field, the Hall conductivity becomes zero. While for the longitudinal conductivity, using the asymptotic expansions

$$\psi(z) = \ln z - \frac{1}{2z} - \frac{1}{12z^2} + \frac{1}{120z^4} + O(\frac{1}{z^5}), \tag{17}$$

we arrive at

$$\begin{aligned}A_L(\omega) &= [\tfrac{(\omega^2-\Gamma^2)}{6\lambda_R^2} - \tfrac{6\lambda_R^2(\omega^2+\Gamma^2)}{(2\Gamma\omega)^2+(6\lambda_R^2)^2}]\ln\tfrac{(-4\lambda_R^2-\omega^2+\Gamma^2)^2+(2\Gamma\omega)^2}{(8\lambda_R^2-\omega^2+\Gamma^2)^2+(2\Gamma\omega)^2} + \tfrac{\Gamma\omega(\omega^2+\Gamma^2)}{(\Gamma\omega)^2+(3\lambda_R^2)^2}[\arctan\tfrac{\omega+2\sqrt{2}\lambda_R}{\Gamma} + \arctan\tfrac{\omega-2\sqrt{2}\lambda_R}{\Gamma}\\ &+ \arctan\tfrac{\omega+2\lambda_R}{\Gamma}\arctan\tfrac{\omega-2\lambda_R}{\Gamma}] - \tfrac{2\Gamma\omega}{3\lambda_R^2}[\arctan\tfrac{\omega+2\sqrt{2}\lambda_R}{\Gamma} + \arctan\tfrac{\omega-2\sqrt{2}\lambda_R}{\Gamma} - \arctan\tfrac{\omega+2\lambda_R}{\Gamma} - \arctan\tfrac{\omega-2\lambda_R}{\Gamma}]\end{aligned} \tag{18}$$

for $\lambda_{SO} = 0$. When $T \to 0$, $|\mu| >> \lambda_R, \Gamma$, the longitudinal conductivity can be further expressed as

$$\sigma_{xx} = \frac{4e^2}{\pi}\frac{|\mu|/\Gamma}{1 + (3\lambda_R^2/\Gamma\mu)^2}. \tag{19}$$

IV. RESULTS AND DISCUSSION

To investigate numerically the behavior of electrical conductivity, we need to restore the whole model parameters in Eqs. (15) and (16). Thus, one should carry out the replacements: $T \to k_BT, eB \to \hbar eBv_F^2$. In the following, we mainly discuss the Rashba SOI effect on the transport properties since the intrinsic SOI is very small, while the Rashba SOI can be tunable by a perpendicular electric field. In Figs. 1 and 2, we show the chemical potential μ dependence of the longitudinal conductivity for the different Rashba spin orbit parameter λ_R at zero or nonzero field. For zero field (see Fig. 1), one can see that when $\lambda_R = 0$, the conductivity is proportional to $|\mu|$ and tends to the known quantum-limited minimal value $4e^2/h$ at zero chemical potential[1]. For $\lambda_R \neq 0$, there exists a threshold chemical potential μ_c which increases with increasing λ_R.



FIG. 1: The longitudinal conductivity σ_{xx} measured in $2e^2/h$ units as a function of the chemical potential μ for the different values of λ_R. We take $B = 0$T, $T = 3$K, $\Gamma = 5$K, and $\lambda_{SO} = 0.001$K.



FIG. 2: The longitudinal conductivity measured in $2e^2/h$ units as a function of the chemical potential μ for the different values of λ_R at $B = 1$T. The other parameters are taken the same as Fig. 1

When the chemical potential is smaller than μ_c, the longitudinal conductivity becomes almost independent of μ; while for $\mu > \mu_c$, the σ_{xx}-μ curves recover the linear relation. This tendency agrees with Eq. (19). For nonzero field case in Fig. 2, we observe SdH oscillations of the conductivity due to the Landau-level crossing of the Fermi level[25]. From Fig. 2, It is clearly seen that when $\lambda_R \neq 0$, each oscillation peak is split into two implicit peaks, and the splitting peaks shift by λ_R. This is due to the spin-orbit splitting of the Landau levels.

The longitudinal conductivity as a function of the magnetic field B for the different λ_R is shown in Fig. 3. For $\lambda_R = 0$, the longitudinal conductivity decreases and intervals between the neighboring SdH oscillation peaks become large with increasing B, which reflects the fact that in the presence of the magnetic field only the transitions between neighboring Landau levels contribute to electrical conductivity, while a further



FIG. 3: The magnetic field dependence of the longitudinal conductivity measured in $2e^2/h$ units for the different values of λ_R at $\mu = -600$K. The other parameters are taken as Fig. 1.

increase of the magnetic field leads to increasing of the distance between neighboring Landau level, thus suppresses the transitions between them. When B is large enough, the conductivity becomes independent of B since the lowest Landau level is filled which is always below the Fermi level. These observations are quite consistent with the previous studies[2526]. In particular, it is interesting to note that when the Rashba SOI presents, the longitudinal conductivity exhibits the characteristic feature that the SdH oscillations are enhanced largely at certain positions, however damped at other positions. Such SdH as a beating pattern have been observed in two dimensional electron gas[27]. From Fig. 3, one can find that the enhanced positions and amplitudes of the SdH oscillations can be tuned by the Rashba SOI due to shift of one set of Landau level by λ_R.



FIG. 4: The longitudinal conductivity measured in $2e^2/h$ units as a function of λ_R for the different magnetic field B at $\mu = -300$K. The other parameters are taken the same as Fig. 1.

Figure 4 shows the longitudinal conductivity versus the Rashba SOI parameter λ_R for the different magnetic field B. It is found that when the magnetic field is applied, the longitudinal conductivity as a function of λ_R behaves as the oscillation. It is because the Rashba SOI leads to the shift of landau level, the longitudinal conductivity shows a maximum each time a Landau level passes through the Fermi level of system, and a minimum when the Fermi level is situated between two Landau levels.



FIG. 5: The Hall conductivity measured in $2e^2/h$ units as a function of the chemical potential μ for the different values of λ_R at $B = 1$T. The other parameters are taken the same as Fig. 1

Figure 5 shows Hall conductivity as a function of the chemical potential μ for the different λ_R. When $\lambda_R = 0$, the Hall conductivity has a steplike structure as a function of μ, which reflect the quantum Hall effect. While the Rashba SOI opens, the Hall steps become narrow and the step near $\mu = 0$ is split into two steps. It is observed that the Hall conductivity displays peaks instead of a plateau at larger λ_R. There is no Hall plateau in the cases of sufficiently strong Rashba SOI. This result suggests that the Rashba SOI tends to suppress the quantum Hall effect in graphene.

V. SUMMARY

We have investigated the effect of the SOIs on transport of Dirac fermions in graphene on the basis of amplitude estimation of Yao *et al.* and Min *et al.* for the SOIs. Using the Schwinger proper-time method, decomposition over Landau level poles and Kubo formula, we obtain the analytical expressions for both longitudinal and Hall conductivities. It has been found that when the Rashba SOI is applied, the longitudinal conductivity versus the chemical potential deviates the linear relation at zero magnetic field. For nonzero magnetic field, the SdH oscillation in the longitudinal conductivity is observed, and each SdH oscillation peak is split into two peaks as the Rashba SOI is applied. While the oscillation in the longitudinal conductivity as a function of the magnetic field exhibits a beating pattern with the Rashba SOI turned on. It is also shown that the Rashba SOI tends to suppress the quantum Hall effect in graphene.

Acknowledgments

Ding gratefully acknowledge financial support from Changsha University of Science and Technology, and Education Department of Hunan Province, China. The work of Zhou was supported by National Natural Science Foundation of China (Grant No. 10574042) and Specialized Research Fund for the Doctoral Program of Higher Education of China (Grant No. 20060542002).

APPENDIX A

The Green functions $G_i^{(R,A)}$ in Eq. (11) are given by

$$\begin{aligned} G_1^{(R,A)}(\omega) &= (H_s + \gamma^0(\omega \pm i\Gamma))[A_1^{(R,A)}(\omega) - i\gamma^1\gamma^2 sgn(eB)A_2^{(R,A)}(\omega)] - i(\gamma^1 k_2 - \gamma^2 k_1)sgn(eB)[B_1^{(R,A)}(\omega) \\ &- A_2^{(R,A)}(\omega)] + (\gamma^1 k_1 + \gamma^2 k_2)[B_2^{(R,A)}(\omega) - A_1^{(R,A)}(\omega)], \end{aligned} \tag{A1}$$

$$\begin{aligned} G_2^{(R,A)}(\omega) &= (H_s + \gamma^0(\omega \pm i\Gamma))\gamma^0 s_z[A_3^{(R,A)}(\omega) - i\gamma^1\gamma^2 sgn(eB)A_4^{(R,A)}(\omega)] - i(\gamma^1 k_2 - \gamma^2 k_1)\gamma^0 s_z sgn(eB)[B_3^{(R,A)}(\omega) \\ &- A_4^{(R,A)}(\omega)] + (\gamma^1 k_1 + \gamma^2 k_2)\gamma^0 s_z[B_4^{(R,A)}(\omega) - A_3^{(R,A)}(\omega)], \end{aligned} \tag{A2}$$

$$\begin{aligned} G_3^{(R,A)}(\omega) &= -\frac{12\lambda_{SO}\lambda_R}{\xi}(H_s + \gamma^0(\omega \pm i\Gamma))(i\gamma^1 s_y - \gamma^0\gamma^1 s_x)(I_4^{(R,A)}(\omega) + \gamma^0 s_z I_8^{(R,A)}(\omega)) + \frac{12\lambda_{SO}\lambda_R}{\xi}(\gamma^1 k_1 \\ &+ \gamma^2 k^2)(I_4^{(R,A)}(\omega) + i\gamma^1\gamma^2 sgn(eB)J_4^{(R,A)}(\omega))(i\gamma^1 s_y - \gamma^0\gamma^1 s_x) + \frac{12\lambda_{SO}\lambda_R}{\xi}(\gamma^1 k_1 + \gamma^2 k^2)(I_8^{(R,A)}(\omega) \\ &+ i\gamma^1\gamma^2 sgn(eB)J_8^{(R,A)}(\omega))(i\gamma^1 s_y - \gamma^0\gamma^1 s_x)\gamma^0 s_z, \end{aligned} \tag{A3}$$

where $A_1 = I_1 + I_3 + I_5 - I_7$, $A_2 = I_2 + I_4/\chi + I_6 - I_8/\chi$, $A_3 = I_1 - I_3 + I_5 + I_7$, $A_4 = I_2 - I_4/\chi + I_6 + I_8/\chi$, $B_1 = I_2 + J_2 + I_6 - J_5$, $B_2 = J_1 + J_3/\chi + J_4 - J_6/\chi$, $B_3 = I_2 - J_2 + I_6 + J_5$, $B_4 = J_1 - J_3/\chi + J_4 + J_6/\chi$, in which

$$I_1^{(R,A)}(\omega) = \tfrac{1}{2}e^{-\alpha}\sum_{n=0}^{\infty}(-1)^n\left[\frac{L_n(2\alpha)-L_{n-1}(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2-\zeta+2n|eB|}+\frac{L_n(2\alpha)-L_{n-1}(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2+\zeta+2n|eB|}\right], \tag{A4}$$

$$I_2^{(R,A)}(\omega) = \tfrac{1}{2}e^{-\alpha}\sum_{n=0}^{\infty}(-1)^n\left[\frac{L_n(2\alpha)+L_{n-1}(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2-\zeta+2n|eB|}+\frac{L_n(2\alpha)+L_{n-1}(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2+\zeta+2n|eB|}\right], \tag{A5}$$

$$\begin{aligned} I_3^{(R,A)}(\omega) = {} & \tfrac{1}{2}e^{-\alpha}\sum_{n=0}^{\infty}(-1)^n\left[\frac{L_n(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2-\zeta-\xi+(2n+1)|eB|}+\frac{L_n(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2+\zeta+\xi+(2n+1)|eB|}\right. \\ & \left.+\frac{L_n(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2-\zeta+\xi+(2n+1)|eB|}+\frac{L_n(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2+\zeta-\xi+(2n+1)|eB|}\right], \end{aligned} \tag{A6}$$

$$\begin{aligned} I_4^{(R,A)}(\omega) = {} & \tfrac{1}{2}e^{-\alpha}\sum_{n=0}^{\infty}(-1)^n\left[\frac{L_n(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2-\zeta-\xi+(2n+1)|eB|}-\frac{L_n(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2+\zeta+\xi+(2n+1)|eB|}\right. \\ & \left.-\frac{L_n(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2-\zeta+\xi+(2n+1)|eB|}+\frac{L_n(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2+\zeta-\xi+(2n+1)|eB|}\right], \end{aligned} \tag{A7}$$

$$I_5^{(R,A)}(\omega) = \tfrac{1}{2}e^{-\alpha}\sum_{n=0}^{\infty}(-1)^n\left[\frac{L_n(2\alpha)-L_{n-1}(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2-\zeta+2n|eB|}-\frac{L_n(2\alpha)-L_{n-1}(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2+\zeta+2n|eB|}\right], \tag{A8}$$

$$I_6^{(R,A)}(\omega) = \tfrac{1}{2}e^{-\alpha}\sum_{n=0}^{\infty}(-1)^n\left[\frac{L_n(2\alpha)+L_{n-1}(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2-\zeta+2n|eB|}-\frac{L_n(2\alpha)+L_{n-1}(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2+\zeta+2n|eB|}\right], \tag{A9}$$

$$\begin{aligned} I_7^{(R,A)}(\omega) = {} & \tfrac{1}{2}e^{-\alpha}\sum_{n=0}^{\infty}(-1)^n\left[\frac{L_n(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2-\zeta-\xi+(2n+1)|eB|}-\frac{L_n(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2+\zeta+\xi+(2n+1)|eB|}\right. \\ & \left.+\frac{L_n(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2-\zeta+\xi+(2n+1)|eB|}-\frac{L_n(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2+\zeta-\xi+(2n+1)|eB|}\right], \end{aligned} \tag{A10}$$

$$\begin{aligned} I_8^{(R,A)}(\omega) = {} & \tfrac{1}{2}e^{-\alpha}\sum_{n=0}^{\infty}(-1)^n\left[\frac{L_n(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2-\zeta-\xi+(2n+1)|eB|}+\frac{L_n(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2+\zeta+\xi+(2n+1)|eB|}\right. \\ & \left.-\frac{L_n(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2-\zeta+\xi+(2n+1)|eB|}-\frac{L_n(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2+\zeta-\xi+(2n+1)|eB|}\right], \end{aligned} \tag{A11}$$

$$J_1^{(R,A)}(\omega) = \tfrac{1}{2}e^{-\alpha}\sum_{n=0}^{\infty}(-1)^n\left[\frac{4L_{n-1}^1(2\alpha)+L_n(2\alpha)-L_{n-1}(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2-\zeta+2n|eB|}+\frac{4L_{n-1}^1(2\alpha)+L_n(2\alpha)-L_{n-1}(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2+\zeta+2n|eB|}\right], \tag{A12}$$

$$\begin{aligned} J_2^{(R,A)}(\omega) = {} & \tfrac{1}{2}e^{-\alpha}\sum_{n=0}^{\infty}(-1)^n\left[\frac{L_n^1(2\alpha)+L_{n-1}^1(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2-\zeta-\xi+(2n+1)|eB|}+\frac{L_n^1(2\alpha)+L_{n-1}^1(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2+\zeta+\xi+(2n+1)|eB|}\right. \\ & \left.+\frac{L_n^1(2\alpha)+L_{n-1}^1(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2-\zeta+\xi+(2n+1)|eB|}+\frac{L_n^1(2\alpha)+L_{n-1}^1(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2+\zeta-\xi+(2n+1)|eB|}\right], \end{aligned} \tag{A13}$$

$$\begin{aligned} J_3^{(R,A)}(\omega) = {} & \tfrac{1}{2}e^{-\alpha}\sum_{n=0}^{\infty}(-1)^n\left[\frac{L_n^1(2\alpha)+L_{n-1}^1(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2-\zeta-\xi+(2n+1)|eB|}-\frac{L_n^1(2\alpha)+L_{n-1}^1(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2+\zeta+\xi+(2n+1)|eB|}\right. \\ & \left.-\frac{L_n^1(2\alpha)+L_{n-1}^1(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2-\zeta+\xi+(2n+1)|eB|}+\frac{L_n^1(2\alpha)+L_{n-1}^1(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2+\zeta-\xi+(2n+1)|eB|}\right], \end{aligned} \tag{A14}$$

$$J_4^{(R,A)}(\omega) = \tfrac{1}{2}e^{-\alpha}\sum_{n=0}^{\infty}(-1)^n\left[\frac{4L_{n-1}^1(2\alpha)+L_n(2\alpha)-L_{n-1}(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2-\zeta+2n|eB|}-\frac{4L_{n-1}^1(2\alpha)+L_n(2\alpha)-L_{n-1}(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2+\zeta+2n|eB|}\right], \tag{A15}$$

$$\begin{aligned} J_5^{(R,A)}(\omega) = {} & \tfrac{1}{2}e^{-\alpha}\sum_{n=0}^{\infty}(-1)^n\left[\frac{L_n^1(2\alpha)+L_{n-1}^1(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2-\zeta-\xi+(2n+1)|eB|}-\frac{L_n^1(2\alpha)+L_{n-1}^1(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2+\zeta+\xi+(2n+1)|eB|}\right. \\ & \left.+\frac{L_n^1(2\alpha)+L_{n-1}^1(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2-\zeta+\xi+(2n+1)|eB|}-\frac{L_n^1(2\alpha)+L_{n-1}^1(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2+\zeta-\xi+(2n+1)|eB|}\right], \end{aligned} \tag{A16}$$

$$\begin{aligned} J_6^{(R,A)}(\omega) = {} & \tfrac{1}{2}e^{-\alpha}\sum_{n=0}^{\infty}(-1)^n\left[\frac{L_n^1(2\alpha)+L_{n-1}^1(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2-\zeta-\xi+(2n+1)|eB|}-\frac{L_n^1(2\alpha)+L_{n-1}^1(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2+\zeta+\xi+(2n+1)|eB|}\right. \\ & \left.+\frac{L_n^1(2\alpha)+L_{n-1}^1(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2-\zeta+\xi+(2n+1)|eB|}-\frac{L_n^1(2\alpha)+L_{n-1}^1(2\alpha)}{(\omega\pm i\Gamma)^2+10\lambda_{SO}^2+2\lambda_R^2+\zeta-\xi+(2n+1)|eB|}\right], \end{aligned} \tag{A17}$$

with $L_n^i(z)$ being the generalized Laguerre polynomials, $\alpha = -\mathbf{k}^2/|eB|$, and $\chi = \xi/|eB|$.

APPENDIX B

Substituting Eq. (11) into the trace Eq. (14), then it is evaluated after a somewhat tedious calculation

$$\begin{aligned}
tr\{\gamma^i S^{(R,A)}(\omega',\mathbf{k})\gamma^j S^{(R,A)}(\omega,\mathbf{k})\} &= \delta_{ij}(\omega' \pm i\Gamma)(\omega \pm i\Gamma)[A_1^{(R,A)}(\omega')A_1^{(R,A)}(\omega) - A_2^{R}(\omega')A_2^{A}(\omega) + A_4^{(R,A)}(\omega')A_4^{(R,A)}(\omega) \\
&- A_3^{(R,A)}(\omega')A_3^{(R,A)}(\omega)] + i\epsilon_{ij}sgn(eB)(\omega' + i\Gamma)(\omega \pm i\Gamma)[A_2^{(R,A)}(\omega')A_1^{(R,A)}(\omega) \\
&- A_1^{(R,A)}(\omega')A_2^{(R,A)}(\omega) + A_3^{(R,A)}(\omega')A_4^{(R,A)}(\omega) - A_4^{(R,A)}(\omega')A_3^{(R,A)}(\omega)] \\
&+ (2k_ik_j - \delta_{ij}\mathbf{k}^2)[(B_1^{(R,A)}(\omega') - A_2^{(R,A)}(\omega'))(B_1^{(R,A)}(\omega) - A_2^{(R,A)}(\omega)) \\
&+ (B_2^{(R,A)}(\omega') - A_1^{(R,A)}(\omega'))(B_2^{(R,A)}(\omega) - A_1^{(R,A)}(\omega)) + (B_3^{(R,A)}(\omega') - A_4^{(R,A)}(\omega')) \\
&\times (B_3^{(R,A)}(\omega) - A_4^{(R,A)}(\omega)) + (B_4^{(R,A)}(\omega') - A_3^{(R,A)}(\omega'))(B_4^{(R,A)}(\omega) - A_3^{(R,A)}(\omega))] \\
&- isgn(eB)[\delta_{ij}(-1)^{i-1}2k_1k_2 + \epsilon_{ij}(k_j^2 - k_i^2)][(B_1^{(R,A)}(\omega') - A_2^{(R,A)}(\omega'))(B_2^{(R,A)}(\omega) - A_1^{(R,A)}(\omega)) \\
&+ (B_3^{(R,A)}(\omega') - A_4^{(R,A)}(\omega'))(B_4^{(R,A)}(\omega) - A_3^{(R,A)}(\omega)) + (B_2^{(R,A)}(\omega') - A_1^{(R,A)}(\omega')) \\
&\times (B_1^{(R,A)}(\omega) - A_2^{(R,A)}(\omega)) + (B_4^{(R,A)}(\omega') - A_3^{(R,A)}(\omega'))(B_3^{(R,A)}(\omega) - A_4^{(R,A)}(\omega))] \\
&+ \delta_{ij}2(\frac{12\lambda_{SO}\lambda_R}{\xi})^2(I_4^{(R,A)}(\omega')I_4^{(R,A)}(\omega) - J_4^{(R,A)}(\omega')J_4 - I_8^{(R,A)}(\omega')I_8^{(R,A)}(\omega) \\
&+ J_8^{(R,A)}(\omega')J_8^{(R,A)}(\omega))\mathbf{k}^2 - i\epsilon_{ij}sgn(eB)2(\frac{12\lambda_{SO}\lambda_R}{\xi})^2[I_4^{(R,A)}(\omega')J_4^{(R,A)}(\omega) \\
&- J_4^{(R,A)}(\omega')I_4^{(R,A)}(\omega) - I_8^{(R,A)}(\omega')J_8^{(R,A)}(\omega) + J_8^{(R,A)}(\omega')I_8^{(R,A)}(\omega)]\mathbf{k}^2, \qquad \text{(B1)}
\end{aligned}$$

where ϵ_{ij} is antisymmetric tensor ($\epsilon_{12} = 1$). Integrating over momenta in Eq. (14), we obtain the longitudinal conductivity

$$\sigma_{xx} = \sigma_{xx}(\Omega \to 0) = \frac{2e^2v_F^2|eB|}{\pi^2}Re\int_{-\infty}^{\infty}d\omega\frac{1}{4T\cosh^2\frac{\beta(\omega-\mu)}{2}}\{X_L + \frac{1}{2|eB|}Y_L - (\frac{6\lambda_{SO}\lambda_R}{\xi})^2Z_L\}, \qquad \text{(B2)}$$

where

$$\begin{aligned}
X_L &= \frac{1}{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi+|eB|}[\frac{2(\omega^2+\Gamma^2)}{i4\Gamma\omega-2\zeta-\xi+|eB|-2|eB|} - \frac{2(\omega+i\Gamma)^2}{-2\zeta-\xi+|eB|-2|eB|}] + \frac{(1-\frac{1}{\chi})}{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi+|eB|} \\
&\times [\frac{(\omega^2+\Gamma^2)}{i4\Gamma\omega-2\zeta+\xi+|eB|-2|eB|} - \frac{(\omega+i\Gamma)^2}{-2\zeta+\xi+|eB|-2|eB|}] - \frac{1}{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta}[\frac{(1-\frac{1}{\chi})(\omega^2+\Gamma^2)}{-i4\Gamma\omega-2\zeta-\xi-|eB|+2|eB|} \\
&- \frac{(1-\frac{1}{\chi})(\omega+i\Gamma)^2}{-2\zeta-\xi-|eB|+2|eB|} + \frac{2(\omega^2+\Gamma^2)}{-i4\Gamma\omega-2\zeta+\xi-|eB|+2|eB|} - \frac{2(\omega+i\Gamma)^2}{-2\zeta+\xi-|eB|+2|eB|}], \qquad \text{(B3)}
\end{aligned}$$

$$\begin{aligned}
Y_L &= w_1\psi(-\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta}{2|eB|}) + w_2\psi(-\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi+|eB|}{2|eB|}) + w_3\psi(-\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi+|eB|}{2|eB|}) \\
&+ w_4\psi(-\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi-|eB|}{2|eB|}) + w_5\psi(-\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi-|eB|}{2|eB|}), \qquad \text{(B4)}
\end{aligned}$$

$$\begin{aligned}
Z_L &= z_1\psi(-\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta+\xi-|eB|}{2|eB|} - z_2\psi(-\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi+|eB|}{2|eB|}) + z_3\psi(-\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi+|eB|}{2|eB|}) \\
&- z_4\psi(-\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta-\xi-|eB|}{2|eB|}) + z_5\psi(-\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi-|eB|}{2|eB|}) - z_6\psi(-\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta+\xi+|eB|}{2|eB|}) \\
&+ z_7\psi(-\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta-\xi+|eB|}{2|eB|}) - z_8\psi(-\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi-|eB|}{2|eB|}), \qquad \text{(B5)}
\end{aligned}$$

with $\psi(z)$ being the digamma function and

$$\begin{aligned}
w_1 &= \frac{2(\omega^2+\Gamma^2)}{-i4\Gamma\omega-2\zeta-\xi+|eB|-2|eB|} - \frac{2(\omega+i\Gamma)^2}{-2\zeta-\xi+|eB|-2|eB|} + \frac{(1-\frac{1}{\chi})(\omega^2+\Gamma^2)}{-i4\Gamma\omega-2\zeta+\xi+|eB|-2|eB|} - \frac{(1-\frac{1}{\chi})(\omega+i\Gamma)^2}{-2\zeta+\xi+|eB|-2|eB|} \\
&+ \frac{(1-\frac{1}{\chi})(\omega^2+\Gamma^2)}{-i4\Gamma\omega-2\zeta-\xi-|eB|+2|eB|} - \frac{(1-\frac{1}{\chi})(\omega+i\Gamma)^2}{-2\zeta-\xi-|eB|+2|eB|} + \frac{2(\omega^2+\Gamma^2)}{-i4\Gamma\omega-2\zeta+\xi-|eB|+2|eB|} - \frac{2(\omega+i\Gamma)^2}{-2\zeta+\xi-|eB|+2|eB|},
\end{aligned}$$

$$w_2 = \frac{2(\omega^2+\Gamma^2)}{-i4\Gamma\omega+2\zeta+\xi-|eB|+2|eB|} - \frac{2(\omega+i\Gamma)^2}{2\zeta+\xi-|eB|+2|eB|}, \quad w_3 = \frac{(1-\frac{1}{\chi})(\omega^2+\Gamma^2)}{-i4\Gamma\omega+2\zeta-\xi-|eB|+2|eB|} - \frac{(1-\frac{1}{\chi})(\omega+i\Gamma)^2}{2\zeta-\xi-|eB|+2|eB|},$$

$$w_4 = \frac{2(\omega^2+\Gamma^2)}{-i4\Gamma\omega+2\zeta-\xi+|eB|-2|eB|} - \frac{2(\omega+i\Gamma)^2}{2\zeta-\xi+|eB|-2|eB|}, \quad w_5 = \frac{(1-\frac{1}{\chi})(\omega^2+\Gamma^2)}{-i4\Gamma\omega+2\zeta+\xi+|eB|-2|eB|} - \frac{(1-\frac{1}{\chi})(\omega+i\Gamma)^2}{2\zeta+\xi+|eB|-2|eB|},$$

$$z_1 = \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta+\xi-|eB|}{-i4\Gamma\omega-2\zeta-2\xi+2|eB|} - \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta+\xi-|eB|}{-i4\Gamma\omega-2\zeta+2|eB|} - \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta+\xi-|eB|}{-2\zeta-2\xi+2|eB|} + \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta+\xi-|eB|}{-2\zeta+2|eB|},$$

$$z_2 = \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi+|eB|}{i4\Gamma\omega-2\zeta-2\xi+2|eB|} - \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi+|eB|}{i4\Gamma\omega-2\zeta+2|eB|} - \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi+|eB|}{-2\zeta-2\xi+2|eB|} + \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi+|eB|}{-2\zeta+2|eB|},$$

$$z_3 = \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi+|eB|}{i4\Gamma\omega-2\zeta+2|eB|} - \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi+|eB|}{i4\Gamma\omega-2\zeta+2\xi+2|eB|} - \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi+|eB|}{-2\zeta+2|eB|} + \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi+|eB|}{-2\zeta+2\xi+2|eB|},$$

$$z_4 = \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta-\xi-|eB|}{-i4\Gamma\omega-2\zeta+2|eB|} - \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta-\xi-|eB|}{-i4\Gamma\omega-2\zeta+2\xi+2|eB|} - \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta-\xi-|eB|}{-2\zeta+2|eB|} + \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta-\xi-|eB|}{-2\zeta+2\xi+2|eB|},$$

$$z_5 = \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi-|eB|}{-i4\Gamma\omega+2\zeta+2\xi+2|eB|} - \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi-|eB|}{-i4\Gamma\omega+2\zeta+2|eB|} - \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi-|eB|}{2\zeta+2\xi+2|eB|} + \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi-|eB|}{2\zeta+2|eB|},$$

$$z_6 = \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta+\xi+|eB|}{i4\Gamma\omega+2\zeta+2\xi+2|eB|} - \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta+\xi+|eB|}{i4\Gamma\omega+2\zeta+2|eB|} - \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta+\xi+|eB|}{2\zeta+2\xi+2|eB|} + \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta+\xi+|eB|}{2\zeta+2|eB|},$$

$$z_7 = \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta-\xi+|eB|}{i4\Gamma\omega+2\zeta+2|eB|} - \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta-\xi+|eB|}{i4\Gamma\omega+2\zeta-2\xi+2|eB|} - \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta-\xi+|eB|}{2\zeta+2|eB|} + \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta-\xi+|eB|}{2\zeta-2\xi+2|eB|},$$

$$z_8 = \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi-|eB|}{-i4\Gamma\omega+2\zeta+2|eB|} - \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi-|eB|}{-i4\Gamma\omega+2\zeta-2\xi+2|eB|} - \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi-|eB|}{2\zeta+2|eB|} + \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi-|eB|}{2\zeta-2\xi+2|eB|}.$$

The Hall conductivity is then given by

$$\sigma_{xy} = \sigma_{xy}(\Omega \to 0) = -\frac{2e^2v_F^2|eB|sgn(eB)}{\pi^2} Im \int_{-\infty}^{\infty} d\omega \frac{1}{4T\cosh^2\frac{\beta(\omega-\mu)}{2}} [X_H + \frac{1}{2|eB|}Y_H + (\frac{6\lambda_{SO}\lambda_R}{\xi})^2 Z_H] \tag{B6}$$

with

$$X_H = a_1/2 - a_1', \quad Y_H = b_1/2 - (b_1' + b_2'), \quad Z_H = c_1/2 - c_1', \tag{B7}$$

where

$$a_1 = \frac{2(1-\frac{1}{\chi})}{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi+|eB|} \frac{\omega^2+\Gamma^2}{-i4\Gamma\omega+2\zeta-\xi-|eB|+2|eB|} + \frac{2(1+\frac{1}{\chi})}{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi+|eB|} \frac{\omega^2+\Gamma^2}{-i4\Gamma\omega+2\zeta+\xi-|eB|+2|eB|} + \frac{\omega^2+\Gamma^2}{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta} [\frac{2(1-\frac{1}{\chi})}{-i4\Gamma\omega-2\zeta-\xi-|eB|+2|eB|} + \frac{2(1+\frac{1}{\chi})}{-i4\Gamma\omega-2\zeta+\xi-|eB|+2|eB|}], \tag{B8}$$

$$b_1 = \mu_1\psi(-\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta}{2|eB|}) - \mu_2\psi(-\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi+|eB|}{2|eB|}) - \mu_3\psi(-\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi+|eB|}{2|eB|}) + \mu_4\psi(-\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi-|eB|}{2|eB|}) + \mu_5\psi(-\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi-|eB|}{2|eB|}), \tag{B9}$$

$$c_1 = \nu_1\psi(-\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta+\xi-|eB|}{2|eB|}) - \nu_2\psi(-\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta-\xi-|eB|}{2|eB|}) + \nu_3\psi(-\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi-|eB|}{2|eB|}) - \nu_4\psi(-\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi-|eB|}{2|eB|}) - \nu_5\psi(-\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta+\xi+|eB|}{2|eB|}) + \nu_6\psi(-\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta-\xi+|eB|}{2|eB|}) - \nu_7\psi(-\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi+|eB|}{2|eB|}) + \nu_8\psi(-\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi+|eB|}{2|eB|}), \tag{B10}$$

$$\begin{aligned} a_1' &= \alpha_0 + \alpha_1 \ln[(\omega+i\Gamma)^2 - (10\lambda_{SO}^2 + 2\lambda_R^2) - \zeta - \xi + |eB|] + \alpha_2 \ln[(\omega+i\Gamma)^2 - (10\lambda_{SO}^2 + 2\lambda_R^2) - \zeta + \xi + |eB|] \\ &+ \alpha_3 \ln[(\omega+i\Gamma)^2 - (10\lambda_{SO}^2 + 2\lambda_R^2) + \zeta], \end{aligned} \tag{B11}$$

$$b_1' = \beta_1 \ln\Gamma(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta}{2|eB|}) + \beta_2\psi(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta}{2|eB|}) + \beta_3\phi(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta}{2|eB|}), \tag{B12}$$

$$\begin{aligned} b_2' &= \beta_4 \ln\Gamma(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi+|eB|}{2|eB|}) + \beta_5\phi(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi+|eB|}{2|eB|}) \\ &- \beta_6\psi(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi+|eB|}{2|eB|}) + \beta_7 \ln\Gamma(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi+|eB|}{2|eB|}) \\ &+ \beta_8\phi(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi+|eB|}{2|eB|}) - \beta_9\psi(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi+|eB|}{2|eB|}) \\ &- \beta_{10} \ln\Gamma(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi-|eB|}{2|eB|}) - \beta_{11}\phi(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi-|eB|}{2|eB|}) \\ &+ \beta_{12}\psi(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi-|eB|}{2|eB|}) - \beta_{13} \ln\Gamma(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi-|eB|}{2|eB|}) \\ &- \beta_{14}\phi(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi-|eB|}{2|eB|}) + \beta_{15}\psi(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi-|eB|}{2|eB|}), \end{aligned} \tag{B13}$$

$$\begin{aligned} c_1' &= \gamma_1 \ln\Gamma(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta+\xi-|eB|}{2|eB|}) + \gamma_4 \ln\Gamma(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta-\xi-|eB|}{2|eB|}) \\ &+ \gamma_7 \ln\Gamma(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi-|eB|}{2|eB|}) + \gamma_{10} \ln\Gamma(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi-|eB|}{2|eB|}) \\ &+ \gamma_{13} \ln\Gamma(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta+\xi+|eB|}{2|eB|}) + \gamma_{16} \ln\Gamma(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta-\xi+|eB|}{2|eB|}) \\ &+ \gamma_{19} \ln\Gamma(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi+|eB|}{2|eB|}) + \gamma_{22} \ln\Gamma(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi+|eB|}{2|eB|}) \\ &+ \gamma_2\psi(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta+\xi-|eB|}{2|eB|}) + \gamma_6\psi(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta-\xi-|eB|}{2|eB|}) \\ &+ \gamma_9\psi(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi-|eB|}{2|eB|}) + \gamma_{12}\psi(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi-|eB|}{2|eB|}) \\ &+ \gamma_{15}\psi(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta+\xi+|eB|}{2|eB|}) + \gamma_{18}\psi(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta-\xi+|eB|}{2|eB|}) \\ &+ \gamma_{21}\psi(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi+|eB|}{2|eB|}) + \gamma_{24}\psi(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi+|eB|}{2|eB|}) \\ &+ \gamma_3\phi(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta+\xi-|eB|}{2|eB|}) + \gamma_5\phi(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta-\xi-|eB|}{2|eB|}) \\ &+ \gamma_8\phi(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi-|eB|}{2|eB|}) + \gamma_{11}\phi(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi-|eB|}{2|eB|}) \\ &+ \gamma_{14}\phi(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta+\xi+|eB|}{2|eB|}) + \gamma_{17}\phi(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta-\xi+|eB|}{2|eB|}) \\ &+ \gamma_{23}\phi(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi+|eB|}{2|eB|}) + \gamma_{20}\phi(-\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi+|eB|}{2|eB|}) \end{aligned} \tag{B14}$$

with $\phi(z) = \int_0^z \ln\Gamma(x)dx$, and

$$\mu_1 = \tfrac{2(1+\frac{1}{\chi})(\omega^2+\Gamma^2)}{-i4\Gamma\omega-2\zeta-\xi+|eB|-2|eB|} - \tfrac{2(1+\frac{1}{\chi})(\omega^2+\Gamma^2)}{-i4\Gamma\omega-2\zeta+\xi-|eB|+2|eB|} + \tfrac{2(1-\frac{1}{\chi})(\omega^2+\Gamma^2)}{-i4\Gamma\omega-2\zeta+\xi+|eB|-2|eB|} - \tfrac{2(1-\frac{1}{\chi})(\omega^2+\Gamma^2)}{-i4\Gamma\omega-2\zeta-\xi-|eB|+2|eB|}, \tag{B15}$$

$$\mu_2 = \tfrac{2(1-\frac{1}{\chi})(\omega^2+\Gamma^2)}{-i4\Gamma\omega+2\zeta-\xi-|eB|+2|eB|}, \quad \mu_3 = \tfrac{2(1+\frac{1}{\chi})(\omega^2+\Gamma^2)}{-i4\Gamma\omega+2\zeta+\xi-|eB|+2|eB|}, \tag{B16}$$

$$\mu_4 = \tfrac{2(1+\frac{1}{\chi})(\omega^2+\Gamma^2)}{-i4\Gamma\omega+2\zeta-\xi+|eB|-2|eB|}, \quad \mu_5 = \tfrac{2(1-\frac{1}{\chi})(\omega^2+\Gamma^2)}{-i4\Gamma\omega+2\zeta+\xi+|eB|-2|eB|}, \tag{B17}$$

$$\nu_1 = 2\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}+2\lambda_R)^2+\zeta+\xi-|eB|}{-i4\Gamma\omega-2\zeta-2\xi+2|eB|} - 2\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}+2\lambda_R)^2+\zeta+\xi-|eB|}{-i4\Gamma\omega-2\zeta+2|eB|}, \tag{B18}$$

$$\nu_2 = 2\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}+2\lambda_R)^2+\zeta-\xi-|eB|}{-i4\Gamma\omega-2\zeta+2|eB|} - 2\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}+2\lambda_R)^2+\zeta-\xi-|eB|}{-i4\Gamma\omega-2\zeta+2\xi+2|eB|}, \tag{B19}$$

$$\nu_3 = 2\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}+2\lambda_R)^2-\zeta-\xi-|eB|}{-i4\Gamma\omega+2\zeta+2\xi+2|eB|} - 2\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}+2\lambda_R)^2-\zeta-\xi-|eB|}{-i4\Gamma\omega+2\zeta+2|eB|}, \tag{B20}$$

$$\nu_4 = 2\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}+2\lambda_R)^2-\zeta+\xi-|eB|}{-i4\Gamma\omega+2\zeta+2|eB|} - 2\tfrac{(\omega+i\Gamma)^2-(10\lambda_{SO}+2\lambda_R)^2-\zeta+\xi-|eB|}{-i4\Gamma\omega+2\zeta-2\xi+2|eB|}, \tag{B21}$$

$$\nu_5 = 2\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}+2\lambda_R)^2+\zeta+\xi+|eB|}{-i4\Gamma\omega-2\zeta-2\xi-2|eB|} - 2\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}+2\lambda_R)^2+\zeta+\xi-|eB|}{-i4\Gamma\omega-2\zeta-2|eB|}, \tag{B22}$$

$$\nu_6 = 2\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}+2\lambda_R)^2+\zeta-\xi+|eB|}{-i4\Gamma\omega-2\zeta-2|eB|} - 2\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}+2\lambda_R)^2+\zeta-\xi+|eB|}{-i4\Gamma\omega-2\zeta+\xi-2|eB|}, \tag{B23}$$

$$\nu_7 = 2\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}+2\lambda_R)^2-\zeta-\xi+|eB|}{-i4\Gamma\omega+2\zeta+2\xi-2|eB|} - 2\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}+2\lambda_R)^2-\zeta-\xi+|eB|}{-i4\Gamma\omega+2\zeta-2|eB|}, \tag{B24}$$

$$\nu_8 = 2\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}+2\lambda_R)^2-\zeta+\xi+|eB|}{-i4\Gamma\omega+2\zeta-2|eB|} - 2\frac{(\omega+i\Gamma)^2-(10\lambda_{SO}+2\lambda_R)^2-\zeta+\xi+|eB|}{-i4\Gamma\omega+2\zeta-2\xi-2|eB|}, \tag{B25}$$

$$\begin{aligned}\alpha_0 &= \frac{2(1+\frac{1}{\chi})(\omega+i\Gamma)^2}{(2\zeta+\xi-|eB|+2|eB|)^2} + \frac{(10\lambda_{SO}^2+2\lambda_R^2)+\zeta+\xi-|eB|}{2\zeta+\xi-|eB|+2|eB|}\frac{\frac{1}{2}(1+\frac{1}{\chi})}{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi+|eB|} + \frac{2(1-\frac{1}{\chi})(\omega+i\Gamma)^2}{(2\zeta-\xi-|eB|+2|eB|)^2} \\ &+ \frac{(10\lambda_{SO}^2+2\lambda_R^2)+\zeta-\xi-|eB|}{2\zeta-\xi-|eB|+2|eB|}\frac{\frac{1}{2}(1-\frac{1}{\chi})}{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi+|eB|} + \frac{2(1+\frac{1}{\chi})(\omega+i\Gamma)^2}{(2\zeta-\xi+|eB|-2|eB|)^2} \\ &+ \frac{(10\lambda_{SO}^2+2\lambda_R^2)-\zeta}{2\zeta-\xi+|eB|-2|eB|}\frac{\frac{1}{2}(1+\frac{1}{\chi})}{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta} + \frac{2(1-\frac{1}{\chi})(\omega+i\Gamma)^2}{(2\zeta+\xi+|eB|-2|eB|)^2} + \frac{(10\lambda_{SO}^2+2\lambda_R^2)-\zeta}{2\zeta+\xi+|eB|-2|eB|}\frac{\frac{1}{2}(1-\frac{1}{\chi})}{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta},\end{aligned}$$

$$\alpha_1 = \frac{2(1+\frac{1}{\chi})[(10\lambda_{SO}^2+2\lambda_R^2)+2\zeta+3\xi-5|eB|]}{(2\zeta+\xi-|eB|+2|eB|)^2}, \quad \alpha_2 = \frac{2(1-\frac{1}{\chi})[(10\lambda_{SO}^2+2\lambda_R^2)+2\zeta-3\xi-5|eB|]}{(2\zeta-\xi-|eB|+2|eB|)^2},$$

$$\alpha_3 = \frac{2(1+\frac{1}{\chi})[(10\lambda_{SO}^2+2\lambda_R^2)-2\zeta-\xi+|eB|-2|eB|]}{(2\zeta-\xi+|eB|-2|eB|)^2} + \frac{2(1-\frac{1}{\chi})[(10\lambda_{SO}^2+2\lambda_R^2)-2\zeta+\xi+|eB|-2|eB|]}{(2\zeta+\xi+|eB|-2|eB|)^2},$$

$$\begin{aligned}\beta_1 &= -\frac{2(1+\frac{1}{\chi})|eB|[2(\omega+i\Gamma)^2+2\zeta+\xi-|eB|+2|eB|]}{(2\zeta+\xi-|eB|+2|eB|)^2} - \frac{2(1-\frac{1}{\chi})|eB|[2(\omega+i\Gamma)^2+2\zeta-\xi-|eB|+2|eB|]}{(2\zeta-\xi-|eB|+2|eB|)^2} \\ &+ \frac{2(1-\frac{1}{\chi})|eB|[2(\omega+i\Gamma)^2+2\zeta+\xi+|eB|-2|eB|]}{(2\zeta+\xi+|eB|-2|eB|)^2} + \frac{2(1+\frac{1}{\chi})|eB|[2(\omega+i\Gamma)^2+2\zeta-\xi+|eB|-2|eB|]}{(2\zeta-\xi+|eB|-2|eB|)^2},\end{aligned}$$

$$\beta_2 = -\frac{(1+\frac{1}{\chi})(\omega+i\Gamma)^2}{2\zeta+\xi-|eB|+2|eB|} - \frac{(1-\frac{1}{\chi})(\omega+i\Gamma)^2}{2\zeta-\xi-|eB|+2|eB|} + \frac{(1-\frac{1}{\chi})(\omega+i\Gamma)^2}{2\zeta+\xi+|eB|-2|eB|} + \frac{(1+\frac{1}{\chi})(\omega+i\Gamma)^2}{2\zeta-\xi+|eB|-2|eB|},$$

$$\beta_3 = -\frac{8|eB|^2(1+\frac{1}{\chi})}{(2\zeta+\xi-|eB|+2|eB|)^2} - \frac{8|eB|^2(1-\frac{1}{\chi})}{(2\zeta-\xi-|eB|+2|eB|)^2} + \frac{8|eB|^2(1-\frac{1}{\chi})}{(2\zeta+\xi+|eB|-2|eB|)^2} + \frac{8|eB|^2(1+\frac{1}{\chi})}{(2\zeta-\xi+|eB|-2|eB|)^2},$$

$$\beta_4 = \frac{2(1+\frac{1}{\chi})|eB|[2(\omega+i\Gamma)^2-2\zeta-\xi+|eB|-2|eB|]}{(2\zeta+\xi-|eB|+2|eB|)^2},$$

$$\beta_5 = \frac{8(1+\frac{1}{\chi})|eB|^2}{(2\zeta+\xi-|eB|+2|eB|)^2}, \quad \beta_6 = \frac{(1+\frac{1}{\chi})(\omega+i\Gamma)^2}{2\zeta+\xi-|eB|+2|eB|},$$

$$\beta_7 = \frac{2(1-\frac{1}{\chi})|eB|[2(\omega+i\Gamma)^2-2\zeta+\xi+|eB|-2|eB|]}{(2\zeta-\xi-|eB|+2|eB|)^2},$$

$$\beta_8 = \frac{8(1-\frac{1}{\chi})|eB|^2}{(2\zeta-\xi-|eB|+2|eB|)^2}, \quad \beta_9 = \frac{(1-\frac{1}{\chi})(\omega+i\Gamma)^2}{2\zeta-\xi-|eB|+2|eB|},$$

$$\beta_{10} = \frac{2(1-\frac{1}{\chi})|eB|[2(\omega+i\Gamma)^2-2\zeta-\xi-|eB|+2|eB|]}{(2\zeta+\xi+|eB|-2|eB|)^2},$$

$$\beta_{11} = \frac{8(1-\frac{1}{\chi})|eB|^2}{(2\zeta+\xi+|eB|-2|eB|)^2}, \quad \beta_{12} = \frac{(1-\frac{1}{\chi})(\omega+i\Gamma)^2}{2\zeta+\xi+|eB|-2|eB|},$$

$$\beta_{13} = \frac{2(1+\frac{1}{\chi})|eB|[2(\omega+i\Gamma)^2-2\zeta+\xi-|eB|+2|eB|]}{(2\zeta-\xi+|eB|-2|eB|)^2},$$

$$\beta_{14} = \frac{8(1+\frac{1}{\lambda})|eB|^2}{(2\zeta-\xi+|eB|-2|eB|)^2}, \quad \beta_{15} = \frac{(1+\frac{1}{\lambda})(\omega+i\Gamma)^2}{2\zeta-\xi+|eB|-2|eB|},$$

$$\gamma_1 = \frac{4|eB|[(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+2\zeta+\xi-2|eB|]}{(-2\zeta+2|eB|)^2} - \frac{4|eB|[(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+2\zeta+2\xi-2|eB|]}{(-2\zeta-2\xi+2|eB|)^2},$$

$$\gamma_2 = \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta+\xi-|eB|}{-2\zeta-2\xi+2|eB|} - \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta+\xi-|eB|}{-2\zeta+2|eB|},$$

$$\gamma_3 = \frac{8|eB|^2}{(-2\zeta+2|eB|)^2} - \frac{8|eB|^2}{(-2\zeta-2\xi+2|eB|)^2},$$

$$\gamma_4 = \frac{4|eB|[(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+2\zeta-\xi-2|eB|]}{(-2\zeta+2|eB|)^2} - \frac{4|eB|[(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+2\zeta-2\xi-2|eB|]}{(-2\zeta+2\xi+2|eB|)^2},$$

$$\gamma_5 = \frac{8|eB|^2}{(-2\zeta+2|eB|)^2} - \frac{8|eB|^2}{(-2\zeta+2\xi+2|eB|)^2},$$

$$\gamma_6 = \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta-\xi-|eB|}{-2\zeta+2\xi+2|eB|} - \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta-\xi-|eB|}{-2\zeta+2|eB|},$$

$$\gamma_7 = \frac{4|eB|[(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-2\zeta-\xi-2|eB|]}{(2\zeta+2|eB|)^2} - \frac{4|eB|[(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-2\zeta-2\xi-2|eB|]}{(2\zeta+2\xi+2|eB|)^2},$$

$$\gamma_8 = \frac{8|eB|^2}{(2\zeta+2|eB|)^2} - \frac{8|eB|^2}{(2\zeta+2\xi+2|eB|)^2},$$

$$\gamma_9 = \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi-|eB|}{2\zeta+2\xi+2|eB|} - \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi-|eB|}{2\zeta+2|eB|},$$

$$\gamma_{10} = \frac{4|eB|[(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-2\zeta+\xi-2|eB|]}{(2\zeta+2|eB|)^2} - \frac{4|eB|[(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-2\zeta+2\xi-2|eB|]}{(2\zeta-2\xi+2|eB|)^2},$$

$$\gamma_{11} = \frac{8|eB|^2}{(2\zeta+2|eB|)^2} - \frac{8|eB|^2}{(2\zeta-2\xi+2|eB|)^2},$$

$$\gamma_{12} = \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi-|eB|}{2\zeta-2\xi+2|eB|} - \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi-|eB|}{2\zeta+2|eB|},$$

$$\gamma_{13} = \frac{4|eB|[(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+2\zeta+2\xi+2|eB|]}{(-2\zeta-2\xi-2|eB|)^2} - \frac{4|eB|[(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+2\zeta+\xi+2|eB|]}{(-2\zeta-2|eB|)^2},$$

$$\gamma_{14} = \frac{8|eB|^2}{(-2\zeta-2\xi-2|eB|)^2} - \frac{8|eB|^2}{(-2\zeta-2|eB|)^2},$$

$$\gamma_{15} = \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta+\xi+|eB|}{-2\zeta-2|eB|} - \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta+\xi+|eB|}{-2\zeta-2\xi-2|eB|},$$

$$\gamma_{16} = \frac{4|eB|[(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+2\zeta-2\xi+2|eB|]}{(-2\zeta+2\xi-2|eB|)^2} - \frac{4|eB|[(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+2\zeta-\xi+2|eB|]}{(-2\zeta-2|eB|)^2},$$

$$\gamma_{17} = \frac{8|eB|^2}{(-2\zeta+2\xi-2|eB|)^2} - \frac{8|eB|^2}{(-2\zeta-2|eB|)^2},$$

$$\gamma_{18} = \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta-\xi+|eB|}{-2\zeta-2|eB|} - \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta-\xi+|eB|}{-2\zeta+2\xi-2|eB|},$$

$$\gamma_{19} = \frac{4|eB|[(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-2\zeta-2\xi+2|eB|]}{(2\zeta+2\xi-2|eB|)^2} - \frac{2|eB|[(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-2\zeta-\xi+2|eB|]}{(2\zeta-2|eB|)^2},$$

$$\gamma_{20} = \frac{8|eB|^2}{(2\zeta+2\xi-2|eB|)^2} - \frac{8|eB|^2}{(2\zeta-2|eB|)^2},$$

$$\gamma_{21} = \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta-\xi+|eB|}{2\zeta-2|eB|} - \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)+\zeta-\xi+|eB|}{2\zeta+2\xi-2|eB|},$$

$$\gamma_{22} = \frac{4|eB|[(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-2\zeta+2\xi+2|eB|]}{(2\zeta-2\xi-2|eB|)^2} - \frac{4|eB|[(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-2\zeta+\xi+2|eB|]}{(2\zeta-2|eB|)^2},$$

$$\gamma_{23} = \frac{8|eB|^2}{(2\zeta-2\xi-2|eB|)^2} - \frac{8|eB|^2}{(2\zeta-2|eB|)^2},$$

$$\gamma_{24} = \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi+|eB|}{2\zeta-2|eB|} - \frac{(\omega+i\Gamma)^2-(10\lambda_{SO}^2+2\lambda_R^2)-\zeta+\xi+|eB|}{2\zeta-2\xi-2|eB|}.$$

The Eqs. (B2) and (B6) are further rewritten as Eqs. (15) and (16).

* Electronic address: dingkaih@mails.gucas.ac.cn
† Electronic address: ghzhou@hunnu.edu.cn

[1] K. S. Novoselov, A. K. Geim, S. V. Morozov, D. Jiang, M. I. Katsnelson, I. V. Grigorieva, S. V. Dubonos, and A. A. Firsov, Nature **438**, 197 (2005).
[2] Y. Zhang, Y. W. Tan, H. L. Stormer, and P. Kim, Nature **438**, 201 (2005).
[3] A. K. Geim and K. S. Novoselov, Nature Materials **6**, 183 (2007).
[4] H. B. Heersche, P. Jarillo-Herrero, J.B. Oostinga, L.M. K. Vandersypen, and A. F. Morpurgo, Nature **446**, 56 (2007).
[5] C. Berger, Z. Song, X. Li, X. Wu, N. Brown, C. Naud, D. Mayou, T. Li, J. Hass, A. N. Marchenkov, E. H.Conrad, P. N. First, and W. A. de Heer, Science **312**, 1191 (2006).
[6] A. H. Castro Neto, F. Guinea, N. M. R. Peres, K. S. Novoselov, A. K. Geim, arXiv:0709.1163.
[7] J. McClure, Phys. Rev. **104**, 666 (1956).
[8] G. W. Semenoff, Phys. Rev. Lett. **53**, 2449 (1984).
[9] F. D. M. Haldane, Phys. Rev. Lett. **61**, 2015 (1988).
[10] K. S. Novoselov, A. K. Geim, S. V. Morozov, D. Jiang, Y. Zhang, S. V. Dubonos, I. V. Grigorieva, and A. A. Firsov, Science **306**, 666 (2004).
[11] K. S. Novoselov, D. Jiang, F. Schedin, T. J.Booth, V. V. Khotkevich, S. V. Morozov, and A. K. Geim, Proc. Natl. Acad. Sci. USA **102**, 10451 (2005).
[12] Y. Zhang, J. P. Small, M. E. S. Amori, and P. Kim, Phys. Rev. Lett. **94**, 176803 (2005).
[13] C. Berger, Z. Song, T. Li, X. Li, A. Y. Ogbazghi, R. Feng, Z. Dai, A. N. Marchenkov, E. H. Conrad, P. N. First, and W. A. de Heer, J. Phys. Chem. B **108**, 19912 (2004).
[14] B. Trauzettel, D. V. Bulaev, D. Loss, and G. Burkard, Nature Physics **3**, 192 (2007).
[15] D. Huertas-Hernando, F. Guinea, and A. Brataas, Phys. Rev. B 74, 155426 (2006).
[16] C. L. Kane and E. J. Mele, Phys. Rev. Lett. **95**, 226801 (2005).
[17] C. L. Kane and E. J. Mele, Phys. Rev. Lett. **95**, 146802 (2005).
[18] Y. Yao, F. Ye, X. L. Qi, S. C. Zhang, and Z. Fang, Phys. Rev. B **75**, 041401(R) (2007).
[19] H. Min, J. E. Hill, N. A. Sinitsyn, B. R. Sahu, L. Kleinman, and A. H. MacDonald, Phys. Rev. B **74**, 165310 (2006).
[20] J. Schwinger, Phys. Rev. **82**, 664 (1951).
[21] A. Chodos, K. Everding, and D. A. Owen Phys. Rev. D **42**, 2881 (1990).
[22] V. P. Gusynin, V. A. Miransky, and I. A. Shovkovy, Phys. Rev. D **52**, 4718 (1995).
[23] G. D. Mahan, *Many-Particle Physics*, Plenum, New York (1990).
[24] Y. Zheng and T. Ando, Phys. Rev. B **65**, 245420 (2002).
[25] S. G. Sharapov, V. P. Gusynin, and H. Beck, Phys. Rev. B **67**, 144509 (2003).
[26] V. P. Gusynin and S. G. Sharapov, Phys. Rev. B **73**, 245411 (2006).
[27] J. Luo, H. Munekata, F. F. Fang, and P. J. Stiles, Phys. Rev. B 41, 7685 (1990).